

LETTER • OPEN ACCESS

# Observed connections of Arctic stratospheric ozone extremes to Northern Hemisphere surface climate

To cite this article: Diane J Ivy *et al* 2017 *Environ. Res. Lett.* **12** 024004

View the [article online](#) for updates and enhancements.

## Related content

- [The surface impacts of Arctic stratospheric ozone anomalies](#)  
K L Smith and L M Polvani
- [On the surface impact of Arctic stratospheric ozone extremes](#)  
N Calvo, L M Polvani and S Solomon
- [A connection from Arctic stratospheric ozone to El Niño-southern oscillation](#)  
Fei Xie, Jianping Li, Wenshou Tian *et al.*

## Recent citations

- [Delayed effect of Arctic stratospheric ozone on tropical rainfall](#)  
Fei Xie *et al*
- [Variations in North Pacific sea surface temperature caused by Arctic stratospheric ozone anomalies](#)  
Fei Xie *et al*



## CEWEG-2017

International Conference on

### Clean Energy for the World's Electricity Grids

United Scientific Group, USA, in collaboration with Fusion Advocates, Geneva, Switzerland is organizing the first International Conference on Clean Energy for the World's Electricity Grids

November 20-22, 2017 Geneva, Switzerland

[click to find out more](#)

[www.ceweg.com](http://www.ceweg.com)

## Environmental Research Letters



## LETTER

## OPEN ACCESS

## RECEIVED

21 July 2016

## REVISED

15 December 2016

## ACCEPTED FOR PUBLICATION

9 January 2017

## PUBLISHED

1 February 2017

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



## Observed connections of Arctic stratospheric ozone extremes to Northern Hemisphere surface climate

Diane J Ivy<sup>1,4</sup>, Susan Solomon<sup>1</sup>, Natalia Calvo<sup>2</sup> and David W J Thompson<sup>3</sup><sup>1</sup> Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139, United States of America<sup>2</sup> Departamento de Física de la Tierra II, Facultad de Ciencias Físicas, Universidad Complutense de Madrid, Madrid, Spain<sup>3</sup> Department of Atmospheric Science, Colorado State University, Fort Collins, CO 80523, United States of America<sup>4</sup> Author to whom any correspondence should be addressed.E-mail: [divy@mit.edu](mailto:divy@mit.edu)**Keywords:** ozone, arctic, tropospheric climateSupplementary material for this article is available [online](#)

## Abstract

We present observational evidence for linkages between extreme Arctic stratospheric ozone anomalies in March and Northern Hemisphere tropospheric climate in spring (March–April). Springs characterized by low Arctic ozone anomalies in March are associated with a stronger, colder polar vortex and circulation anomalies consistent with the positive polarity of the Northern Annular Mode/North Atlantic Oscillation in March and April. The associated spring tropospheric circulation anomalies indicate a poleward shift of zonal winds at 500 hPa over the North Atlantic. Furthermore, correlations between March Arctic ozone and March–April surface temperatures reveal certain regions where a surprisingly large fraction of the interannual variability in spring surface temperatures is associated with interannual variability in ozone. We also find that years with low March Arctic ozone in the stratosphere display surface maximum daily temperatures in March–April that are colder than normal over southeastern Europe and southern Asia, but warmer than normal over northern Asia, adding to the warming from increasing well-mixed greenhouse gases in those locations. The results shown here do not establish causality, but nevertheless suggest that March stratospheric ozone is a useful indicator of spring averaged (March–April) tropospheric climate in certain Northern Hemispheric regions.

## 1. Introduction

The pronounced stratospheric ozone depletion in the Southern Hemisphere (known as the ozone hole) forms under extreme cold temperatures in the lower stratosphere over the Antarctic and has been an annual feature in austral spring since its discovery in the 1980s (Farman *et al* 1985). The Antarctic ozone hole has resulted in a colder and stronger polar vortex and a positive trend in the Southern Annular Mode (SAM), which has influenced Southern Hemispheric surface climate in austral summer (Thompson and Solomon 2002, Gillett and Thompson 2003, Thompson *et al* 2011). In contrast, Arctic ozone losses are generally much weaker and more variable than in the Antarctic (Solomon *et al* 2014). The Arctic winter circulation and temperatures are more variable due to enhanced planetary wave driving in the Northern Hemisphere.

Dynamically active winters typically experience major sudden stratospheric warmings, when the zonal-mean circulation reverses from westerly to easterly at 10 hPa, which can warm the Arctic stratosphere by up to tens of degrees (Charlton and Polvani 2007) and limit stratospheric ozone losses. Consequently, dynamically active winters are associated with positive stratospheric ozone anomalies and circulation anomalies consistent with the negative polarity of the Northern Annular Mode (NAM). In comparison, dynamically quiescent Arctic winters since the late 1970s are associated with colder temperatures and negative ozone anomalies, due to both enhanced chemical depletion and weakened transport (Shaw and Perlwitz 2014), along with circulation anomalies consistent with the positive polarity of the NAM/North Atlantic Oscillation (NAO).

While the relationship between the ozone hole at stratospheric altitudes and Southern Hemispheric

summer surface climate has been supported by multiple studies (Thompson and Solomon 2002, Previdi and Polvani 2014 and references therein), the observed relationships between Arctic stratospheric ozone and Northern Hemispheric extratropical circulation are less clear. However, it is known that ozone anomalies are related to circulation anomalies in the stratosphere, and previous studies have focused on the relationship between stratospheric circulation anomalies and tropospheric climate. Circulation anomalies in the Northern Hemispheric extratropical upper stratosphere appear to propagate down into the lower stratosphere, where they may persist for several weeks, and influence both the mean tropospheric circulation and the incidence of extreme weather events in winter (Baldwin and Dunkerton 2001, Thompson and Wallace 2001). In particular, winters with major sudden stratospheric warmings, which are associated with circulation anomalies resembling the negative polarity of the NAM, show surface pressure anomalies resembling the negative polarity of the NAO, an increase in blocking over Greenland and Iceland, and a decrease in blocking over the Eastern Atlantic and Europe (Lehtonen and Karpechko 2016, Davini *et al* 2014). Moreover, the tropospheric response following stratospheric final warmings (when the polar vortex breaks down for the last time in spring) also resemble the transition of the NAO from positive to negative polarity (Black *et al* 2006).

Observational evidence to date has not yet indicated a causal linkage between Arctic ozone and tropospheric climate. Recent modeling studies have examined the possible connections using a range of approaches, and obtained mixed results. One study by Cheung *et al* (2014) used the 2011 extreme Arctic ozone anomalies reported by the Earth Observing System (EOS) Microwave Limb Sounder (MLS) to probe whether the extreme Arctic ozone depletion of 2011 had an effect on the prediction of tropospheric climate. They found no improvement in spring tropospheric forecast skill between simulations with the UK Met Office operational weather forecasting model. Karpechko *et al* (2014) found a relationship between the 2011 low Arctic stratospheric ozone anomalies with tropospheric climate in ECHAM5 simulations with specified sea surface temperatures, but noted that specifying the ozone anomalies alone did not result in a significant surface impact. In contrast, Smith and Polvani (2014) used an atmospheric global climate model to study whether extreme low Arctic ozone anomalies could affect Northern Hemispheric climate. Analysis of time-slice experiments revealed a colder, stronger Arctic polar vortex in their simulations using prescribed ozone anomalies based upon total ozone data. The prescribed ozone forcing needed for a robust tropospheric response in their simulations appeared to be larger than that historically observed. A coupled chemistry-climate simulation by Calvo *et al* (2015) found a robust


stratospheric-tropospheric response in low versus high ozone years: a poleward shift of the North Atlantic tropospheric jet, a positive phase of the NAO, and a corresponding response in surface temperatures in late spring. Their study used an ensemble of simulations driven by historically observed ozone depleting substances. The fully-coupled approach of Calvo *et al* (2015) allows consistency between the evolving ozone distributions and dynamical conditions, which may explain the differences between their conclusions and those of studies prescribing fixed ozone concentrations. However, whether differences in the ozone forcings between the various studies could contribute to the range of conclusions has not been systematically evaluated.

In this study, we analyze observational datasets to further explore the connection between high versus low spring Arctic ozone (hereafter referred to as differences) and subsequent Northern Hemispheric climate. We examine the relationship of Arctic ozone differences in March to the tropospheric circulation and surface climate in spring averages (March–April). Baldwin and Dunkerton (2001) characterize the stratospheric behavior as a ‘harbinger’ of tropospheric change—one that need not indicate causality but can be useful as an indicator; we take the same approach here. Ozone is readily measured with very high precision on a routine basis, motivating our examination of its utility as a possible climatic indicator. As both increasing well-mixed greenhouse gases and El Niño–Southern Oscillation (ENSO) also affect Northern Hemispheric climate through stratospheric pathways (Butler *et al* 2014, Calvo *et al* 2011), we examine their impacts on stratospheric and tropospheric climate as well, and evaluate possible overlaps with ozone. In addition, since low/high ozone anomalies are generally observed in dynamically quiescent/active winters, we analyze winters with and without major sudden stratospheric warmings for comparison.

## 2. Methods

### 2.1. Datasets

The differences in Arctic stratospheric and Northern Hemispheric extratropical tropospheric climate for high versus low ozone years are calculated here using a variety of datasets. Vertically resolved temperature, wind, ozone, geopotential height, and mean sea level pressure were mainly taken from monthly-mean reanalysis data from the National Aeronautics and Space Administration’s Modern-Era Retrospective Analysis for Research and Applications (MERRA, Rienecker *et al* 2011) dataset. Data from MERRA are available since the satellite era began in 1979 and at a horizontal resolution of  $0.5^\circ$  latitude by  $0.67^\circ$  longitude and a vertical resolution of 42 levels that extends from the surface to 0.1 hPa. The reanalysis data from the Interim European Centre for Medium-Range



Weather Forecasts (ERA-Interim, Dee *et al* 2011) are also used; ERA-Interim provides data since 1979 at 1° latitude by 1° longitude and 37 vertical levels from the surface to 1 hPa. The monthly-mean surface temperatures used here are based on ERA-Interim as well as monthly-mean and daily station data from the National Climatic Data Center's Global Historical Climatology Network (GHCNV3, Lawrimore *et al* 2011). The near-surface winds (925 hPa) are zonal and meridional monthly-mean reanalysis data from ERA-Interim.

## 2.2. Analysis

Changes in stratospheric and tropospheric climate were estimated as the differences between subsets of years determined by the potential forcing being analyzed, where the potential forcings include: 1) extreme low and high Arctic ozone anomalies, 2) occurrence of a major sudden stratospheric warming, 3) ENSO events, and 4) increasing well-mixed greenhouse gases.

Based on the limited available record, springs with extreme ozone anomalies were identified as the seven years with the highest and seven years with the lowest ozone abundances from 1979–2012 for both total column and vertically resolved lower stratospheric (70 hPa) polar cap averaged (63–90°N) ozone from

MERRA as shown in figure 1; for brevity we present the tropospheric responses using the vertically resolved lower stratospheric Arctic ozone. MERRA's total column ozone is primarily based on SBUV measurements, with TOMS/OMI data used to fill in missing SBUV data. MERRA's vertically resolved ozone is estimated by MERRA's Data Assimilation System (DAS) from transported constituents of the odd oxygen family and includes constraints to SBUV partial and total column measurements and an ozone climatology from McPeters *et al* (2007) as a prior.

Statistical significance between the subsets of data was evaluated using a Student's *t* statistic (as in Santer *et al* 2000) and are presented at the 95% confidence level based on a one-sided test. Pearson correlation coefficients are also calculated between linearly detrended March ozone and linearly detrended spring (March–April) surface temperatures and winds using monthly- or seasonal-mean data from 1979–2012. The statistical significance of the correlation coefficient was also evaluated using a Student's *t* statistic. Significant changes in daily maximum spring temperatures for selected regions are determined from probability distributions of daily GHCN station data. The distributions were estimated assuming a bin size of 20, estimated from the square root of the number of



observations available in the selected set of years for Southern Asia (which had the most limited record); statistical significance was determined using a two-sample Kolmogorov-Smirnov test.

In this study, we chose March to represent spring Arctic ozone, which is the month with the largest observed ozone losses and is consistent with the constructed ozone forcings used in Smith and Polvani (2014). Calvo *et al* (2015) selected April as the indicator month; our analysis suggests that the observed maxima in ozone depletion peak one month earlier on average than in their model. We analyze the tropospheric response over March and April (hereafter referred to as spring). The tropospheric responses in March and April are fairly similar (discussed further in the results section). Therefore, we present the tropospheric responses in the spring average, to provide an improved signal to noise and for comparison with the earlier modeling studies.

Years with major sudden stratospheric warmings were identified using the climatologies from Charlton and Polvani (2007), for 1958–2002, and Kuttippurath and Nikulin (2012), for 2003–2012. Both studies use the criterion of zonal-mean zonal wind reversal from westerly to easterly at 60°N at 10 hPa to identify the onset of a major sudden stratospheric warming. Years with a strong influence from El Niño or La Niña were identified as when the National Weather Service Climate Prediction Center's Oceanic Niño Index (ONI) exceeded  $\pm 1$  in any month from November through March ([www.cpc.ncep.noaa.gov/products/analysis\\_monitoring/ensostuff/ensoyears.shtml](http://www.cpc.ncep.noaa.gov/products/analysis_monitoring/ensostuff/ensoyears.shtml)). The ONI is calculated as the three month running mean of sea surface temperature anomalies from ERSSTv4 over the Niño-3.4 region (5°S–5°N, 120°W–170°W), where the anomalies are estimated from a 30-year period, which is updated every 5 years. The role of increasing well-mixed greenhouse gases was estimated as the difference between the years 2007–2012 and 1979–1984 (referred to below as 'historical').

### 3. Results

Figure 1 shows the seasonal cycle and interannual variability of total column and lower stratospheric (70 hPa) ozone from MERRA averaged over the Arctic. Years with low ozone anomalies in March are characterized by lower ozone abundances in early winter as well, and these anomalies persist throughout the summer. Figure 1 also shows time series of March total column and lower stratospheric ozone at 70 hPa with the ozone extremes (as defined in section 2.2) highlighted. Winters with major sudden stratospheric warmings and strong La Niña or El Niño events are also indicated. Low ozone anomalies predominate in the late 1990s. High ozone anomalies are observed in both the early 1980s as well as in the 2000s, which helps to distinguish the relationship of tropospheric climate


with ozone from that of increasing well-mixed greenhouse gases.

Randel *et al* (2009) noted higher Northern Hemispheric polar ozone abundances during El Niño winters, as El Niño winters have a weaker polar vortex due to enhanced planetary wave activity (e.g. van Loon and Labitzke 1987). Iza and Calvo (2015) also found that the polar vortex was weaker in El Niño winters not only during canonical El Niño (Eastern Pacific El Niño) events but also during Central Pacific El Niño events coinciding with major sudden stratospheric warmings. In contrast, La Niña winters are characterized by a stronger vortex and negative ozone anomalies (Garfinkel and Hartmann 2007, Free and Seidel 2009). However, Butler and Polvani (2011) showed that the frequency of sudden stratospheric warming events was similar in El Niño and La Niña winters. As seen in figure 1, lower ozone abundances are overall observed in winters without a major sudden stratospheric warming, as compared to winters with them, while the relationship between extremes in Arctic spring ozone at 70 hPa and ENSO is less clear. This is consistent with Rieder *et al* (2013), who found that the fingerprint of ENSO on ozone was strongest in the tropics and mid-latitudes and diminished over the polar regions.

To illustrate the tropospheric impact of stratospheric ozone anomalies, figure 2 (leftmost column) presents the seasonal differences in polar cap (75–90°N) averaged ozone, temperature, and geopotential height, as well as zonal winds averaged over the North Atlantic (60°W–0°E) from 55–65°N between years with low March ozone compared to high March ozone anomalies. Years characterized by low ozone anomalies in March show lower ozone throughout winter and into the fall in the lower stratosphere (left column of figure 2), as also seen in figure 1. Colder temperatures are also observed in late winter and persist through the fall in the lower stratosphere in years with low ozone.

Lower geopotential heights are observed in late winter and into the summer in the stratosphere in years with low ozone, consistent with the positive polarity of the NAM/NAO and corresponding to a stronger, colder polar vortex in the stratosphere and a poleward shift of the mid-latitude jet in the troposphere. Over the midlatitudes, years with low ozone anomalies show stronger North Atlantic zonal winds in winter and spring (bottom left panel of figure 2).

The additional columns of figure 2 show the behavior obtained in the same variables when different potential forcings are considered: years without major sudden stratospheric warmings compared to years with them; years with strong ENSO events, and the long-term change. Figure 2 shows that that the strongest and most robust differences in Arctic stratospheric and tropospheric winter and spring climate occur for the differences between years with



low March ozone anomalies compared to high ozone anomalies. Differences in Arctic climate between years without major sudden stratospheric warmings and with them show similar patterns to the differences based on low and high March ozone, albeit weaker. This is not surprising as years with low/high March ozone anomalies tend to occur in winters without/with major sudden stratospheric warmings (figure 1). Figure S1 (available at [stacks.iop.org/ERL/12/024004/mmedia](https://stacks.iop.org/ERL/12/024004/mmedia)) shows that the temperature changes obtained in figure 2 using MERRA are similar to those found using radiosonde station data from HadAT2, version 2 (Thorne *et al* 2005); therefore, these changes are not artifacts of the reanalysis.


Overall the stratospheric-tropospheric differences shown in figure 2 between La Niña and El Niño years are much weaker than those associated with interannual variability of ozone and the occurrence of major sudden stratospheric warmings. Note that Calvo *et al* (2011) found that only El Niño winters with SSWs have a robust impact on tropospheric climate through the stratosphere. The historical differences between the early 1980s and late 2010s in the Arctic stratospheric climate (right column of figure 2) show a weaker, warmer polar vortex in early winter, followed

by a colder polar vortex in late spring, albeit weaker than in the analysis with ozone extremes. However, studies have shown that historical trends or differences in Arctic winter stratospheric climate are sensitive to the chosen end-year, suggesting that the historical responses are not robust to the chosen end-year (Manney *et al* 2005, Ivy *et al* 2014).

Figure 2 indicates that interannual variations of ozone in March project onto zonal-mean surface climate in spring. Figure S2 further explores the correlations between March ozone at 70 hPa with zonally-varying zonal wind at 500 hPa, sea level pressure, and surface temperatures in March, April, and the spring average (March and April). As seen in figure S2, there are regions with significant correlations between the Northern Hemisphere tropospheric climate and March ozone in both March and April individually, and the patterns exhibit several similar features in the two months, particularly in the surface temperature field (figure S2 bottom row). The tropospheric changes project onto the zonal-mean (figure 2) but also display strong zonal asymmetries (figure S2).


Figure 3 shows maps summarizing the regional differences in Northern Hemispheric tropospheric climate in spring between years with low and high





ozone abundances in March. The figure is very similar to the right column of figure S2, but not identical since it is based on composite differences rather than correlations using March ozone (see also figure 4 in Smith and Polvani (2014) and figure 3 in Calvo *et al* (2015)). Figure S3 compares the results in figure 3 with the attendant results calculated for major sudden

stratospheric warmings, ENSO, and the long-term trend. Years with low ozone are associated with a statistically significant poleward shift of the midlatitude jet over the North Atlantic (figure 3 top) that is similar to that noted in previous modeling studies (Smith and Polvani 2014, Calvo *et al* 2015). The differences in sea level pressure are consistent with



the positive polarity of the NAM/NAO, with lower sea level pressures over the Arctic and higher pressures over the midlatitudes in spring for years with low ozone anomalies. The differences in spring surface temperatures show warmer temperatures over northern Siberia and southwestern Europe, and colder temperatures over southeastern Europe and southern Asia in years with low March ozone anomalies (figure 3).

To highlight the relationships between March ozone and spring surface temperatures on interannual timescales, figure 4 shows the correlation coefficients of March Arctic ozone at 70 hPa with spring averaged surface temperatures and near-surface winds from ERA-Interim. The temperature results are similar to those in the composite difference analysis of figure 3 and identical to those in correlation analysis in the right column of figure S2. Over southeastern Europe

and southern Asia, years with low ozone anomalies tend to have colder surface temperatures, as seen by the positive correlations. Over northern Asia, low ozone anomalies are associated with warmer surface temperatures, corresponding to southwesterly winds bringing warmer air poleward (as seen in the wind vectors in figure 4). Overall, more areas are statistically significant than in the composite analysis due to the inclusion of more years in the analysis. The associated time series of surface temperatures for selected locations are shown in figure 4 (middle and bottom panels) and illustrate locally strong relationships with March Arctic spring ozone, with correlations of 0.46 for Grand Rapids, MI, USA,  $-0.67$  for eastern Siberia,  $-0.77$  for central Russia, and  $0.67$  for Nepal; all are statistically significant at the 95% level.

To investigate the possible relationship between spring ozone and daily surface temperature extremes,






figure 5 shows the probability distributions for daily maximum temperatures in spring for selected regions displaying high correlation coefficients between March ozone and linearly detrended spring surface temperatures (figure 4). The distributions of daily maximum temperature between the high and low ozone years in the regions shown in figure 5 were all statistically different at the 95% level based on a two-sample Kolmogorov-Smirnov test. Figure 5 (right) also shows that some of the regions show strong differences in the historical time period, as would be expected from increasing well-mixed greenhouse gases (as also seen in figure S3). Over southeastern Europe and southern Asia, years with low ozone anomalies display colder extreme spring temperatures that appear to be more extreme than those in the historical time period. Over Siberia and central Asia, years characterized by low ozone show warmer temperatures than years with high ozone, but also reveal substantial changes in the historical time period. Thus, figure 5 suggests that the connection between stratospheric ozone changes on extreme surface temperature may be comparable to those associated with increasing GHGs over certain regions.

#### 4. Conclusions

Previous studies (Baldwin and Dunkerton 2001, Thompson and Wallace 2001) have shown direct

coupling between the stratospheric and tropospheric circulations in NH winter. Our analysis illustrates the value of ozone measurements as an indicator for stratospheric circulation change, and shows that the tropospheric anomalies persist after the extreme ozone anomalies for up to one month. Springs characterized by low ozone anomalies in March show a stronger, colder polar vortex in the stratosphere in late winter and early spring. In the troposphere, years with negative ozone anomalies during March are associated with circulation anomalies consistent with the positive polarity of the NAM/NAO that persist into April, including a poleward shift of the North Atlantic jet, lower than normal temperatures over eastern North America, southeastern Europe, and southern Asia, and higher than normal temperatures over northern and central Asia. Furthermore, we suggest that these stratospheric-tropospheric coupled results in spring are most pronounced when analyzed in the context of Arctic ozone extremes, and generally less apparent when analyzed for ENSO events or increasing greenhouse gases.

Our analysis of several observational datasets supports a linkage between March Arctic lower stratospheric ozone and spring Northern Hemispheric extratropical surface climate. Although there is a limited number of extreme ozone years available for analysis, broad agreement between the observations and modeling results gives confidence in our findings

(Calvo *et al* 2015). Further, correlations between March stratospheric ozone and surface temperatures averaged for March and April are significant in certain locations even when all years are considered (figure 4). Finally, while it is clear that tropospheric greenhouse gas increases have driven many changes in surface climate extremes in recent decades, this study suggests that stratospheric extremes are significant contributors to extreme surface temperatures in spring at some locations (figure 5). While the linkages between ozone and surface climate highlighted here do not necessarily imply causality, our results do indicate predictability of March ozone for spring tropospheric climate, particularly in certain regions. Future work is needed to evaluate the predictive skill of using ozone for Northern Hemispheric tropospheric climate.

## Acknowledgments

We thank two anonymous reviewers for their helpful comments and suggestions. DJI and SS were supported by National Science Foundation (NSF) Atmospheric Chemistry grant AGS-1539972. NC acknowledges partial support from the Spanish Ministry of Economy and Competitiveness through the PALEOSTRAT project (CGL2015-69699) project and the European Project 603557-STRATOCLIM under program FP7-ENV.2013.6.1-2. DWJT is supported by the NSF Climate Dynamics Program.

## References

- Baldwin M P and Dunkerton T J 2001 Stratospheric harbingers of anomalous weather regimes *Science* **294** 581–84
- Barnes E A, Dunn-Sigouin E, Masato G and Woollings T 2014 Exploring recent trends in Northern Hemisphere blocking *Geophys. Res. Lett.* **41** 638–44
- Barriopedro D and Calvo N 2014 On the relationship between ENSO, stratospheric sudden warmings, and blocking *J. Clim.* **27** 4704–20
- Black R X, McDaniel B A and Robinson W A 2006 Stratosphere-troposphere coupling during spring onset *J. Clim.* **19** 4891–901
- Butler A H and Polvani L M 2011 El Niño, La Niña, and stratospheric sudden warmings: A reevaluation in light of the observational record *Geophys. Res. Lett.* **38** L13807
- Butler A H, Polvani L M and Deser C 2014 Separating the stratospheric and tropospheric pathways of El Niño–Southern Oscillation teleconnections *Environ. Res. Lett.* **9** 024014
- Calvo N and Marsh D R 2011 The combined effects of ENSO and the 11 year solar cycle on the Northern Hemisphere polar stratosphere *J. Geophys. Res.* **116** D23112
- Calvo N, Polvani L M and Solomon S 2015 On the surface impact of Arctic stratospheric ozone extremes *Environ. Res. Lett.* **10** 094003
- Charlton A J and Polvani L M 2007 A new look at stratospheric sudden warmings. Part I: Climatology and modeling benchmarks *J. Clim.* **20** 449–69
- Cheung J C H, Haigh J D and Jackson D R 2014 Impact of EOS MLS ozone data on medium-extended range ensemble weather forecasts *J. Geophys. Res. Atmos.* **119** 9253–66
- Davini P, Cagnazzo C and Anstey J A 2014 A blocking view of the stratosphere-troposphere coupling *J. Geophys. Res. Atmos.* **119** 100–115
- Dee D P *et al* 2011 The ERA-Interim reanalysis: configuration and performance of the data assimilation system *Q.J.R. Meteorol. Soc.* **137** 553–97
- Farman J C, Gardiner B G and Shanklin J D 1985 Large losses of total ozone in Antarctica reveal seasonal ClO<sub>x</sub>/NO<sub>x</sub> interaction *Nature* **315** 207–10
- Free M and Seidel D J 2009 Observed El Niño–Southern Oscillation temperature signal in the stratosphere *J. Geophys. Res.* **114** D23108
- Garfinkel C I and Hartmann D L 2007 Effects of the El Niño Southern Oscillation and the Quasi-Biennial Oscillation on polar temperatures in the stratosphere *J. Geophys. Res.* **112** D19112
- Gillet N P and Thompson D W J 2003 Simulation of recent Southern Hemispheric climate change *Science* **302** 273–5
- Hardiman S C *et al* 2011 Improved predictability of the troposphere using stratospheric final warmings *J. Geophys. Res.* **116** D18113
- Ivy D J, Solomon S and Thompson D W J 2014 On the identification of the downward propagation of Arctic stratospheric climate change over recent decades *J. Clim.* **27** 2789–99
- Iza M and Calvo N 2015 Role of stratospheric sudden warmings on the response to central pacific El Niño *Geophys. Res. Lett.* **42** 2482–9
- Karpechko A Y, Perlwitz J and Manzini E 2014 A model study of tropospheric impacts of the Arctic ozone depletion of 2011 *J. Geophys. Res.* **119** 7999–8014
- Kuttippurath J and Nikulin G 2012 A comparative study of the major sudden stratospheric warmings in the Arctic winters 2003/2004–2009/2010 *Atmos. Chem. Phys.* **12** 8115–29
- Lawrimore J H, Menne M J, Gleason B E, Williams C N, Wuertz D B, Vose R S and Rennie J 2011 An overview of the Global Historical Climatology Network monthly mean temperature data set, version 3 *J. Geophys. Res.* **116** D19121
- Lehtonen I and Karpechko A Y 2016 Observed and modelled tropospheric cold anomalies associated with sudden stratospheric warmings *J. Geophys. Res. Atmos.* **121** 1591–610
- Manney G L, Kruger K, Sabutis J L, Sena S A and Pawson S 2005 The remarkable 2003–2004 winter and other recent warm winters in the Arctic stratosphere since the late 1990s *J. Geophys. Res.* **110** D04107
- McPeters R D, Labow G J and Logan J A 2007 Ozone climatological profiles for satellite retrieval algorithms *J. Geophys. Res.* **112** D05308
- Previdi M and Polvani L 2014 Climate system response to stratospheric ozone depletion and recovery *Q. J. Roy. Met. Soc.* **140** 2401–19
- Randel W J, Garcia R R, Calvo N and Marsh D 2009 ENSO influence on zonal mean temperature and ozone in the tropical lower stratosphere *Geophys. Res. Lett.* **36** L15822
- Renwick J A and Wallace J M 1996 Relationships between North Pacific Wintertime Blocking, El Niño, and the PNA pattern *Mon. Weather Rev.* **124** 2071–6
- Rieder H E, Frossard L, Ribatet M, Staehelin J, Maeder J A, Di Rocco S, Davison A C, Peter T, Weihs P and Holawe F 2013 On the relationship between total ozone and atmospheric dynamics and chemistry at mid-latitudes—Part 2: The effects of the El Niño/Southern Oscillation, volcanic eruptions and contributions of atmospheric dynamics and chemistry to long-term total ozone changes *Atmos. Chem. Phys.* **13** 165–79
- Rienecker M *et al* 2011 MERRA: NASA's Modern-Era Retrospective analysis for Research and Applications *J. Clim.* **24** 3624–48
- Santer B D, Wigley T M L, Boyle J S, Gaffen D J, Hnilo J J, Nychka D, Parker D E and Taylor K E 2000 Statistical significance of trends and trend differences in layer-average atmospheric temperature time series *J. Geophys. Res. Atmos.* **105** 7337–56

- Shaw T A and Perlwitz J 2014 On the control of the residual circulation and stratospheric temperatures in the Arctic by planetary wave coupling *J. Atmos. Sci.* **71** 195–206
- Smith K L and Polvani L M 2014 The surface impacts of Arctic stratospheric ozone anomalies *Environ. Res. Lett.* **9** 074015
- Solomon S 1999 Stratospheric ozone depletion: A review of concepts and history *Rev. Geophys.* **37** 275–316
- Solomon S, Haskins J, Ivy D J and Min F 2014 Fundamental differences between Arctic and Antarctic ozone depletion *Proc. Natl Acad. Sci.* **111** 6220–25
- Thompson D W J and Wallace J M 2001 Regional climate impacts of the Northern Hemisphere Annular Mode *Science* **293** 85–9
- Thompson D W J and Solomon S 2002 Interpretation of recent Southern Hemisphere climate change *Science* **296** 895–99
- Thompson D W J, Solomon S, Kushner P J, England M H, Grise K M and Karoly D J 2011 Signatures of the Antarctic ozone hole in Southern Hemisphere surface climate change *Nat. Geosc.* **4** 741–9
- Thorne P W, Parker D E, Tett S F B, Jones P D, McCarthy M, Coleman H and Brohan P 2005 Revisiting radiosonde upper air temperatures from 1958 to 2002 *J. Geophys. Res.* **110** D18105
- Uppala S M, Kallberg P W, Simmons A J *et al* 2005 The ERA-40 Reanalysis *Q.J.R. Meteorol. Soc.* **131** 2961–3012
- van Loon H and Labitzke K 1987 The Southern Oscillation. Part V: The anomalies in the lower stratosphere of the Northern Hemisphere in winter and a comparison with the Quasi-Biennial Oscillation *Mon. Weather Rev.* **115** 357–69